

## A REVIEW OF THE MEDICINAL FERNS OF UKRAINE

Valentyna Minarchenko<sup>1</sup>, Iryna Tymchenko<sup>2</sup>, Tetyana Dvirna<sup>2</sup>, Larysa Makhynia<sup>1</sup>

<sup>1</sup>*Department of Pharmacognosy and Botany,  
Bohomolets National Medical University, Kyiv, Ukraine*

<sup>2</sup>*Department of Systematics and Floristics of Vascular Plants,  
M.G. Kholodny Institute of Botany, National Academy of Sciences of Ukraine*

### ABSTRACT

The growing demand for natural medicines, food products and cosmetics stimulates the search and studies on the diversity of plant species. Ferns are a promising source of biologically active substances for pharmaceutical and medical purposes. Out of 63 species of wild ferns in Ukraine, 39 species contain biologically active substances, which are used, or can be used for medicinal and food purposes. Most of them have limited distribution and resources. Many of them are endangered and are in need of protection. This article highlights some aspects of the distribution, resource significance, and protection status in Ukraine and/or Europe, as well as the medicinal value of the wild medicinal ferns of Ukraine.

**Keywords:** *wild medicinal ferns, resources significance, protection*

### INTRODUCTION

Today the investigation of the available diversity and resources of medicinal plants has become an important subject and is progressing in the spheres of biodiversity and conservation. Ferns are an important component of phytodiversity and have value as medicinal, ornamental, edible and technical plants in different regions of the world.

The ferns are thought by most people to have low value for medical purposes. In the traditional medicine of Ukraine and that of most of the European countries, their useful aspects are largely limited. Ferns are popular plants as food, medicine, orna-

mental, fiber, crafts and building material mostly in the countries of Southeast Asia (1-6).

Nowadays, the medicinal properties of ferns in regard to human health are being more thoroughly studied. According to various data, ferns contain many secondary metabolites, as well as polyphenols, flavonoids, triterpenoids, etc. (7-8). Ferns have various functional effects on human health, such as antioxidant, antimicrobial, antitumor, anti-inflammatory and more, which are classified as the effects of secondary metabolites (9-10). It has been affirmed that antioxidant activity of the secondary metabolites of ferns is the most useful bioactivity for countering aging and chronic disease (1).

Therefore, this study was conducted in order to create a database of wild medicinal ferns of Ukraine, their resource significance, protection status in Ukraine or Europe and medicinal value, based on a long-term research, results of literature data analysis and local ethnobotanical knowledge. There are nearly 63 wild species, subspecies and hybrids of fern in Ukraine (11). In this study, a total of 39 wild species of ferns native to Ukraine, which are considered medicinal in the world, were analyzed. This research

---

#### Address for correspondence:

Valentyna Minarchenko  
Department of Pharmacognosy and Botany  
Bohomolets National Medical University  
13 Tarasa Shevchenko Blvd  
01601 Kyiv  
Ukraine  
e-mail: valminar@ukr.net

**Received:** May 2, 2017

**Accepted:** June 27, 2017

Table 1. The state and medicinal values of ferns of Ukraine

| Latin name                            | English name           | Family                         | Distribution in Ukraine | Re-source value | Number of individuals in population | Protection status in Ukraine | Protection status (by Global and European regional assessment) | Medicinal value                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|---------------------------------------|------------------------|--------------------------------|-------------------------|-----------------|-------------------------------------|------------------------------|----------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>Adiantum capillus-veneris</i> L.   | Venus hair fern        | Adiantaceae (Pteridaceae s.l.) | L                       | 0               | F                                   | RB, endangered               | LC                                                             | Flavonoids, triterpenoids, phenylpropanoids, carbohydrates, carotenoids. The presence of bitterness explains the use of leaves as a vermifuge. Used as expectorant, antitussive, febrifuge, galactagogue, emollient, antiparasitic and dandruff, to increase lactation, weak emmenagogue and kidney function, weak tonic, sudorific, emetic and astrigent effect; anti-inflammatory, antiviral, antimicrobial, antifungal and significant analgesic activity (15, 16); used by women who had just given birth in order to restore menstrual blood. It is anemollient, and adjuvant in bronchial diseases (17-19). |
| <i>Asplenium adiantum-nigrum</i> L.   | Black spleenwort       | Aspleniaceae                   | R                       | 0               | F                                   | RB, rare                     |                                                                | Saponins, triterpenoids, higher fatty acids, higher aliphatic carbohydrates, phenolcarboxylic acids and their derivatives, xanthones (mahniferin); inflammatory, laxative, lactogenic, anthelmintic, diuretic, expectorant, diseases of the spleen and jaundice (6, 20); in experiment - anesthetic, strong antioxidant and hepatoprotective activity (21); in diseases of the spleen, jaundice, conjunctivitis (4, 7).                                                                                                                                                                                           |
| <i>Asplenium cuneifolium</i> Viv.     | Serpentine spleenwort  | Aspleniaceae                   | L                       | 0               | I                                   | RP                           |                                                                | Phytoestrogens, lipophilic substances; tocopherol, cholesterol, sitosterol, stigma sterols; estrogenic activity, in traditional medicine to treat menstrual pain and as a contraceptive (22).                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <i>Asplenium fontanum</i> (L.) Bernh. | Smooth rock spleenwort | Aspleniaceae                   | L                       | 0               | I                                   |                              |                                                                | Kaempferol 3-gentiobioside, kaempferol 3-polyglycoside, kaempferol 3,7-glycoside (24-25).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |

|                                                                     |                       |                   |   |   |   |    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|---------------------------------------------------------------------|-----------------------|-------------------|---|---|---|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Asplenium<br>ruta-muraria<br>L.                                     | Wall rue              | Asplenia-<br>ceae | M | 0 | F | RP | Caffeic acid glycoside, 2-O-caffeoyl- $\beta$ -D-fructofuranosyl-(2 $\rightarrow$ 1)- $\alpha$ -D-glucopyranoside and an ( $\alpha$ , $\beta$ )-isomeric pair of 2E-caffeoyl-D-glucopyranoside, together with kaempferol-3-O- $\beta$ -D-[6-E-caffeoyl- $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 2) glucopyranoside]-7-O- $\beta$ -D-glucopyranoside, 1-O-caffeoyl glycoside, sucrose, diploptene and $\beta$ -sitosterol (26, 27); astringent, diuretic, anthelmintic, expectorant; for respiratory diseases, respiratory infections, jaundice, ascites, scurvy (7).                                                                                                                                                                         |
| Asplenium<br>scolopendrium L. (Phyllitis scolopendrium (L.) Newman) | Hart's<br>tongue fern | Asplenia-<br>ceae | M | I | I | RP | Organic acids, triterpenoids, steroids, nitrogen-containing compounds, lipids, higher aliphatic hydrocarbons, flavonoids, flavonol glycosides (7, 28-30); astringent, laxative, analgesic, wound healing, hemostatic, detoxification, hemostatic and expectorant effect. Externally it is used as an ointment in the treatment of piles, burns and scalds. An infusion is taken internally for the treatment of diarrhea, dysentery, gravelly deposits of the bladder and for removing obstructions of the liver and spleen; strong antioxidant activities (5); in folk medicine it is used for tuberculosis, infertility, diseases and tumors of the stomach, liver, spleen; in homeopathy in cases of pulmonary tuberculosis and malaria (8). |
| Asplenium<br>septentrionale<br>(L.) Hoffm.                          | Forked<br>spleenwort  | Asplenia-<br>ceae | M | 0 | F | RP | Organic acids, nitrogen-containing compounds, terpenoids, flavonoids (kaempferol, quercetin) (29); aldehydes, alcohols, especially fatty acids are abundant (43%); sesquiterpene hydrocarbons and oxygenated diterpene phytol (31). Astringent, expectorant; liver diseases and respiratory diseases (4, 7).                                                                                                                                                                                                                                                                                                                                                                                                                                    |

|                                 |                         |              |   |     |   |    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|---------------------------------|-------------------------|--------------|---|-----|---|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Asplenium trichomanes L.        | Maidenhair spleenwort   | Aspleniaceae | M | 0   | F | RP | Triterpenoids, nitrogen-containing compounds, flavonoids (kaempferol, quercetin) (31); polyketide and aromatic compounds, isoprenoid derivatives (33); laxative, anthelmintic, diuretic, expectorant, sedative, abortive, estrogenic activity (22, 34); in case of bronchitis, respiratory infections, depression, jaundice, ascites.<br>For irregular menses, breast diseases, coughs, and liver ailments (7, 8).                                                                                                                                                    |
| Asplenium viride Huds.          | Green spleenwort        | Aspleniaceae | R | 0   | I | RP | Flavonoids: kaempferol, quercetin (35). Powder from the leaves of the plant is applied externally for injuries and wounds. As tea – for gastric ulcer gastritis (36, 37).                                                                                                                                                                                                                                                                                                                                                                                             |
| Athyrium filix-femina (L.) Roth | Lady-fern (Female fern) | Athyriaceae  | M | III | N | RP | Triterpenoids and polyphenols, carotenoids, higher fatty acids, steroids, tannins, vitamin C (7, 38); cytotoxic, antioxidant and antimicrobial activity (5); analgesic, astringent, expectorant, weak anthelmintic and diuretic; use in the treatment of pregnant women to prevent their water from breaking prematurely, for aiding mothers with intestinal fevers, to help with breast pain, caked breasts and for treating men with venereal diseases. For treatment enterocolitis, dysentery, nephritis, metrorrhagia, hemorrhoids, headaches and rheumatism (7). |
| Azolla filiculoides Lam.        | Water fern              | Azollaceae   | R | I   | N |    | Flavonoids (anthocyanidins: luteolinidin and apigenin); a mixture of lipids, unsaturated lipids, polysaccharides, polyphenols (O-dihydroxyphenols with free OH groups and tannins) and alkaloids; hepatoprotective and antioxidant activity (39–42).                                                                                                                                                                                                                                                                                                                  |
| Azolla mexicana C. Presl        | Mexican mosquito-fern   | Azollaceae   | R | I   | F |    | Flavonoids (43).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |

|                                                                       |                                                      |                                          |   |   |   |                   |                                                                                                                                                                                                                                                                                                                                                                                                                          |
|-----------------------------------------------------------------------|------------------------------------------------------|------------------------------------------|---|---|---|-------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Blechnum<br>spicant (L.)<br>Roth                                      | Hard fern                                            | Blechnaceae                              | R | 0 | I | RP                | <p>Triterpenoids, flavonoids, steroids, higher aliphatic alcohols; in tumors of the spleen; slight antithrombin activity or wound healing (7).</p> <p>The leaflets in folk medicine have been chewed in the treatment of internal cancer, lung disorders and stomach problems. The fronds are used externally as a medicine for skin sores. A decoction of the root has been used in the treatment of diarrhea (44).</p> |
| Botrychium<br>lunaria (L.)<br>Sw.                                     | Common<br>moonwort                                   | Ophioglossaceae (incl.<br>Botrychiaceae) | R | 0 | I | RB,<br>vulnerable | <p>Moonwort is broadly antibacterial against many microbes, including most fungus and some viruses; used as a good vulnerary and also used in dysentery (45); it can be used on various kinds of injuries, pulled muscles, insect stings, contusions, cuts, infections, nerve pain and in anything itchy and irritated like contact dermatitis (7); also used in the treatment of hernias and dysentery.</p>             |
| Botrychium<br>matricariifolium (A. Braun<br>ex Döll) W. D.<br>J. Koch | Chamomile<br>grape-fern<br>(Matricary<br>grape-fern) | Ophioglossaceae (incl.<br>Botrychiaceae) | R | 0 | I | NT                | <p>Leaves used as wound healing (7).</p>                                                                                                                                                                                                                                                                                                                                                                                 |
| Botrychium<br>multifidum (S.G. Gmel.)<br>Rupr.                        | Leathery<br>grape-fern,<br>(Leathery<br>moonwort)    | Ophioglossaceae (incl.<br>Botrychiaceae) | R | 0 | I | DD                | <p>Leaves used as wound healing (7).</p>                                                                                                                                                                                                                                                                                                                                                                                 |
| Ceterach<br>officinarium<br>Willd.                                    | Rusty-back<br>fern                                   | Aspleniaceae                             | L | 0 | I |                   | <p>Triterpenoids, phenolcarboxylic acids, flavonoids, carbohydrates, tannins, higher fatty acid; used in diseases and tumors of the spleen, kidneys, bladder, skin diseases (7, 46); as an astringent and anti-inflammatory on reddened or irritated skin, applied using poultices of crushed or fresh leaves. It is also known as an expectorant (47).</p>                                                              |
| Cheilanthes<br>acrosticha<br>(Balb.) Tod.                             | Fragrant<br>cheilanthes                              |                                          | L | 0 | I | RB,<br>endangered | <p>Carbohydrates, flavonoids (7), used in toothache (48).</p>                                                                                                                                                                                                                                                                                                                                                            |

|                                                                                      |                                             |                  |   |    |   |          |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|--------------------------------------------------------------------------------------|---------------------------------------------|------------------|---|----|---|----------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>Cystopteris fragilis</i> (L.) Bernh.                                              | Bladder fern                                | Cystopteridaceae | L | 0  | I | RB, rare | Proteins, carbohydrates, phenolic compounds, terpenoids (4); xanthones, phenols and their derivatives; toning, softening, expectorant, antipyretic, anthelmintic, bacteriostatic, analgesic; intestinal diseases, asthma, physical fatigue (7), treating trauma (44).                                                                                                                                                                                                                                                                 |
| <i>Dryopteris carthusiana</i> (Vill.) H.P.Fuchs                                      | Narrow buckler fern                         | Dryopteridaceae  | M | II | N | RP       | Para-aspidin BB. Rhizomes can be used as an anthelmintic. Water and alcohol extracts of rhizomes, water extract of leaves show bacteriostatic activity (9). In Belarus rhizomes are used for dermatomycosis (7).                                                                                                                                                                                                                                                                                                                      |
| <i>Dryopteris caucasica</i> (A. Braun) Fraser-Jenk. & Corley                         | Caucasian buckler fern                      | Dryopteridaceae  | L | I  | I |          | Phloroglucines: dezaspidin BP, para-aspidin BB, PB (9).                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| <i>Dryopteris cristata</i> (L.) A. Gray                                              | Crested buckler fern                        | Dryopteridaceae  | M | I  | N | RP       | Para-aspidin BB, PB, desaspidin BP, flavaspidic acid PB, flavaspidic acid PP (9); anthelmintic, expectorant, bacteriostatic (7).                                                                                                                                                                                                                                                                                                                                                                                                      |
| <i>Dryopteris dilatata</i> (Hoffm.) A. Gray                                          | Broad buckler fern                          | Dryopteridaceae  | M | I  | F | RP       | Astragalin, isoquercitrin and chlorogenic acid, caffeoyltartrate and kaempferol-3-O- $\beta$ -D-glucoside-7-O-a-L-rhamnoside were detected (49). Polyketide and aromatic compounds, isoprenoid derivative (33), desaspidin BP, flavaspidic acid BB (9). The root contains 'flicin', a substance that paralyzes tapeworms and other internal parasites and has been used as a worm expellent. It is one of the most effective treatments known for tapeworms. An infusion of the fronds is used as a hair rinse and to treat dandruff. |
| <i>Dryopteris expansa</i> (c. Presl) Fraser-Jenkis & Jermy (=D. assimilis S. Walker) | Alpine buckler fern (Northern buckler fern) | Dryopteridaceae  | R | 0  | I | RP       | Phenols and their derivatives desaspidin BB, tris-desaspidin BBB and phloropyron BB (9); anthocyanins, essential oils, saponins; anthelmintic, analgesic, wound healing, antibacterial (7).                                                                                                                                                                                                                                                                                                                                           |

|                                                                                                |                    |                  |   |     |   |    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|------------------------------------------------------------------------------------------------|--------------------|------------------|---|-----|---|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Dryopteris filix-mas (L.) Schott                                                               | Male fern          | Dryopteridaceae  | M | III | N | RP | Astragal, isoquercitrin and chlorogenic acid, rutin and kaempferol-3-O-rutinoside (49) Para-aspidin, flavaspidic acid PB, filixic acid BBB (9). Major compounds the phloroglucinols, have broad spectrum of pharmacological activities e.g. anthelmintic or antirheumatic effects (49). Rhizomes are used in folk medicine of many countries as a remedy against tapeworms. In Russia a drug "Phylisan" (Filixanum) from the rhizome Dryopteris filix-mas was used as anthelmintic remedy (7).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Dryopteris villarii (Bellardi) Woytn. ex Schinz & Thell. (=Dryopteris mindshelkensis N. Pavl.) | Rigid buckler-fern | Dryopteridaceae  | L | I   | F |    | Phloroglucines: albaspidin AP, AB, PP, PB, BB, apigenin 7-O-glucoside-4'-acetate, quercetin 3-O-rhamnoside-7-O-glucoside (9, 50).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Gymnocarpium dryopteris (L.) Newman                                                            | Common oak fern    | Cystopteridaceae | M | I   | F | RP | Carbohydrates, proteins, free amino acids, saponins, phenolic compounds, tannins, volatile oils, terpenoids (4). Possessing anti-inflammatory, anthelmintic, analgesic, antiseptic and other useful properties, it is known in folk medicine of many European and Asian countries. The flavonol glycoside astragal possesses anti-inflammatory, anti-allergenic and antimicrobial activities. It together with other flavonoids is used to treat rheumatoid arthritis and osteoarthritis (51). Tincture of the leaves, as well as decoctions from the leaves, is used as an analgesic and anti-inflammatory agent for rheumatic pains, arthralgia, arthritis and arthrosis. From the decoction of leaves lotions for the treatment of wounds, ulcers, mastitis, bruises are made. The aqueous decoction of fern roots has an anthelmintic effect. It is used for diseases of the spleen, certain diseases of the digestive tract and excretory system (nephritis, cystitis) (7). |

|                                                                          |                                      |                         |   |   |   |                |    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|--------------------------------------------------------------------------|--------------------------------------|-------------------------|---|---|---|----------------|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Marsilea quadrifolia L.                                                  | Water shamrock (Common water clover) | Marsileaceae            | R | 0 | I | RB, vulnerable | NT | Carbohydrates, alkaloids, steroids, tannins, terpenes, protein, flavonoids and anthraquinone. Juice made from the leaves is diuretic and febrifuge and also used to treat snake bite and applied to abscesses etc. The plant is anti-inflammatory, diuretic, depurative, febrifuge, refrigerant (6); has antibacterial cytotoxic, antioxidant and anticancer activity. The plants is useful in vitiated pitta, cough, bronchitis, diabetes, psychiatric diseases, eye diseases, diarrhea and skin disease (52). It has got profound antibacterial, cytotoxic and antioxidant effect and may have potential use in medicine (53).                                                 |
| Onoclea struthiopteris (L.) Hoffm. (Matteuccia struthiopteris (L.) Tod.) | Ostrich fern                         | Onocleaceae             | M | 0 | F | RP             | LC | A caffeic acid derivate, flavonoids, stilbenes and phenolic components: demethoxymatteucinol, matteucinol, matteuorin, pinosylvin and pinosylvin 3-O- $\beta$ -Dglucopyranoside, 5- $\beta$ -Dglucosyloxy-3-hydroxyl-trans-stilbene-2-carboxylic acid (54-59). The rhizomes and frond bases have been used as a traditional Chinese medicine for the treatment of pinworms, influenza, dysentery, hematochezia and uterine hemorrhage. The pharmacological activities of plants include antiviral, antiparasitic activities, and eliminating bacteria. Rhizomes were used for heavy menstruation, coughing, asphyxiation, as an astringent, fever and malignant tumors (60, 61). |
| Ophioglossum vulgatum L.                                                 | Adder's fern                         | Ophioglossaceae (incl.) | M | I | F | RP             |    | Flavonoids (quercetin, kaempferol) (7). Wound healing and tissue repair properties. Traditionally used on wounds and burns in the form of ointments (62, 63).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Oreopteris limbosperma (Bellardi & All.) Holub                           | Lemon-scented fern                   | Thelypteridaceae        | L | 0 | I | RP             |    | Terpenic compounds incl. (E)-nerolidol, alpha-terpineol, beta-caryophyllene, linalool, pinenes, limonene, and gamma-terpinen-7-al) and carotenoid-type derivatives (64).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Pilularia globulifera L.                                                 | Pillwort                             | Marsileaceae            | L | 0 | I | RB, endangered | NT | Flavonoids (quercetin 3-O-rhamnoside, kaempferol) (7, 65).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |

|                                 |                                |                 |   |   |   |    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|---------------------------------|--------------------------------|-----------------|---|---|---|----|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Polypodium vulgare L.           | Common polypody (Wall fern)    | Polypodiaceae   | M | I | F | RP | LC | <p>Triterpenoids, steroids, carbohydrates, organic acids, phenols and their derivatives, essential oils (sesquiterpenes (39.6%) and their oxidized forms (21.9%), fatty oils, catechins (31). Polypody rhizome contains: ecdysteroids, phloroglucinol derivatives, volatile oil, 8% fixed oil and tannins. Polypody root contains also sugar, saponins, glycyrrhizin, mannitol, protein, starch and carbohydrates (slime) and calcium malate (66). The data of clinical pharmacology is not available, but the antibacterial effect was investigated (67). It is used in traditional medicine as an expectorant in cough and cold, and constipation. It stimulates bile secretion and is a gentle laxative. In European herbal medicine it is traditionally used as a treatment for hepatitis and jaundice and as a remedy for indigestion and loss of appetite. It should not be used externally since it can cause skin rashes (44, 66).</p> <p>It has previously been used as a remedy against respiratory complaints and rheumatism. It has expectorant and laxative effect.</p> |
| Polystichum aculeatum (L.) Roth | Hard shield-fern               | Dryopteridaceae | R | I | I | RP |    | <p>Triterpenoids, cardiac glycosides, polyphenols, anthraquinones and quinines; anthelmintic, antibacterial, strong antioxidant activity (5, 7).</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Polystichum lonchitis (L.) Roth | Alpine rough fern (Holly fern) | Dryopteridaceae | R | 0 | I | RP |    | <p>Lipids (spores); cholagogic, wound healing, diuretic properties; used in diseases of the spleen, skin rash (7, 68).</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |

|                                                                            |                                    |                  |   |     |   |                    |    |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|----------------------------------------------------------------------------|------------------------------------|------------------|---|-----|---|--------------------|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Pteridium aquilinum (L.) Kuhn (incl. Pteridium tauricum Krecz. ex Grossh.) | Common bracken                     | Dennstaedtiaceae | M | III | N |                    |    | Organic acids, carotenoids, sesquiterpenes and norsesquiterpenes, steroids, flavonoids, phenolic compounds, starches, tannins, lipids; anti-diabetic and anti-obesity activities, antioxidant, diuretic, astringent, antipyretic, antitussive, anthelmintic, healing, bacteriostatic (69). In Chinese medicine - diuretic, antipyretic, with infectious hepatitis; in Indian medicine - in spleen infiltrate. In folk medicine - laxative, tonic, wound healing; in diseases of the respiratory, stomachalgia, myalgia, diarrhea (7, 16). Young leaves - used as a food in East Asia and South Africa. The use of its rhizomes and fronds as food appears to be widespread. Young croziers are consumed fresh, canned, dried or pickled, and they are regarded as a delicacy in many countries. However, consumption of the bracken fern has been shown to induce bladder and intestinal carcinomas in cattle and to cause a number of diseases in other farm animals. Some human populations also eat young bracken shoots and epidemiological studies in Japan and Brazil have shown a close association between bracken consumption and the occurrence of certain cancers. An unstable glucoside named ptaquiloside has been proven to have potent carcinogenicity (70). |
| Salvinia natans (L.) All.                                                  | Floating moss (Floating watermoss) | Salviniaceae     | M | 0   | N | RB, data deficient | LC | Glycosides, carbohydrates, proteins, tannins, terpenoids, flavonoids, phospholipids, fatty acids are the main classes of compounds was isolated from the leaves of <i>Salvinia natans</i> (4). Antioxidant properties were investigated (28, 71). In traditional Chinese medicine it is used as an antipyretic, analgesic, against poison, increasing circulation of blood (72).                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |

|                                                                              |                |                  |   |    |   |                |    |                                                                                                                                                              |
|------------------------------------------------------------------------------|----------------|------------------|---|----|---|----------------|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Thelypteris palustris Schott (=Thelypteris confluentis (Thunb.) C.V. Morton) | Marsh fern     | Thelypteridaceae | M | II | N | RP             | LC | Flavonoid compounds (73). Used in gynecology and as an anthelmintic agent (68).                                                                              |
| Woodsia ilvensis (L.) R.Br.                                                  | Oblong woodsia | Woodsiaceae      | L | 0  | I | RB, endangered |    | Essential oils: monoterpenes, diterpenes, sesquiterpenes, alcohols, aldehydes, acids (31); tannins; antipyretic, wound healing; for kidney diseases (7, 68). |

ABBREVIATIONS

Distribution in Ukraine: M- on much of the territory, R- in some regions, L - known in some locations;  
Resources value: III- the raw resources of wild species are sufficient for harvesting in large volume, II- resources are limited, I- resources are not available, species needs to be protected ;  
Number of representatives in a population: N - numerous, F - few in numbers, I - isolated individuals;  
Protection status in Ukraine: RB - Red Book, RP - regional Protection;  
Protection status (by Global and European regional assessment): NT - Near Threatened, LC - Least Concern, DD - Data Deficient.

is part of a complex study of the medicinal plants of Ukraine with a goal to provide necessary information about the diversity, raw resources and necessity of protection of medicinal plants in the state.

MATERIALS AND METHODS

Regular field studies (expeditions) were carried out in different regions of Ukraine during the years 2002-2016. In the field, we have made geobotanical descriptions of localities, studied medicinal species and performed analysis of their involvement in plant communities. Additional information about the diversity, distribution, protection status, medicinal properties and use of ferns was received as a result of analyzing herbarium materials from various institutions of Ukraine, population survey during trips and literature data.

RESULTS AND DISCUSSION

Ferns do not hold significant economic importance in Ukraine. They are mostly used for decorative purposes in landscape gardening and occasionally in folk medicine for health benefits . Some of them (*Pteridium aquilinum* (L.) Kuhn, *Onoclea struthiopteris* (L.) Hoffm. and *Matteuccia struthiopteris* (L.) Tod.) are occasionally used for cooking purposes. It is a good source of protein and trace elements.

The results of the survey are presented in Table 1, in which the plants are arranged in alphabetical order. For each species the following data is present: : Latin and English name of fern species, name of family, and family, distribution and resource significance in Ukraine, population structure, protection status in Ukraine and Europe, and medicinal value.

The presented species of ferns belong to 16 families, with the obvious predominance of the *Aspleniaceae* (23%) and *Dryopteridaceae* (23%) families. Other families are few in numbers and mostly represented by small quantity of species.

A distribution analysis shows widespread species, which grow on most of the territory of Ukraine (41.02%). The rest of the ferns have a limited distribution in some regions (30.77%) (Polissya, Carpathian Mountains, forest-steppe zone, etc.) or are only known to be in certain locations (28.21%). The last of these are rare ones and included to Red Book of Ukraine (12). Their populations are small with few representatives. Most of the fern species of Ukraine

being shade and moisture-loving, grow under forest canopy or in specific habitats, mainly in the Polissya region. These forests are being heavily transformed by logging. Any disturbance of forest vegetation leads to ecological imbalance and destruction of most of the fern populations. With the increase of environment transformation the natural habitats of ferns are becoming small and fragmented, so some species are on the brink of extinction and protected at regional or national levels (see Table 1).

Based on the results of the evaluation of the resource potential of the analyzed ferns, they were divided into several groups of species (see Table 1). Based on estimation, only *Athyrium filix-femina* (L.) Roth, *Dryopteris filix-mas* (L.) Schott and *Pteridium aquilinum* have resource significance and are available for sustainable wild collection and wide economic importance. The resources of other species are limited and inaccessible for harvesting from the wild. Most of them are protected on regional, national or global level.

Thus, more than half of the medicinal ferns have little distribution in Ukraine and many of them are endangered and are in need of protection. Their populations, mostly, are small-numbered or represented by isolated individuals. For this reason, 11 species of medicinal ferns from the total number studied are protected at the state (national) level and are included in the Red Book of Ukraine (12); 19 species are protected at regional level and listed on the Regional Rare Checklists. Some of them are protected only in certain areas (administrative regions), where there is a threat of depletion of their populations. Besides that, ten species from the total number studied are threatened at an European level (see Table 1), according to data from global or European regional assessment (13).

The protection rank at European level of *Botrychium matricariifolium* (A. Braun ex Döll) W.D.J.Koch, *Marsilea quadrifolia* L. and *Pilularia globulifera* L. has been identified as Near Threatened. There are rare species with a scattered distribution throughout Europe. They are listed as threatened on all available national red lists, including Ukraine's, and populations are declining almost everywhere. Species, such as *Adiantum capillus-veneris* L., *Botrychium lunaria* (L.) Sw., *Salvinia natans*

(L.) All., *Onoclea struthiopteris*, *Polypodium vulgare* L. and *Thelypteris palustris* (Salib.) Schott are classified in the European Red List as Least Concern as they are widespread and abundant, however they appear to be declining throughout much of their range with some localities going extinct (13). The first three species are added to the Red Book of Ukraine, however the populations of *Salvinia natans* have been found lately in many localities with a high number of representatives and the overall trend is stable with range extensions around some river basins. *Onoclea struthiopteris*, *Polypodium vulgare* and *Thelypteris palustris* are abundant in large areas, although protected in some regions. *Botrychium multifidum* (S.G. Gmel.) Rupr. has been identified as Data Deficient (DD) on European level, but is in decline in Ukraine and included in the Red Book as rare species (Table 1). In addition, the Red Book of Ukraine includes *Asplenium adiantum-nigrum* L., *Cheilanthes acrostica* (Balb.) Tod., *Cystopteris fragilis* (L.) Bernh. and *Woodsia ilvensis* (L.) R.Br., as populations with a decreasing number of representatives.

The current main threat for many wild ferns of Ukraine is disturbance of the ecological balance of their living environment, including: general ecosystem dehumidification, wetland amelioration, forest exploitation (deforestation, intensive logging, decrease of the area of old-growth forests), habitat fragmentation as a result of anthropogenic transformation of the environment, wild fern collection, livestock grazing and related agricultural threats, local fires, recreation and tourism. The most sensitive to these threats are stenotopic species, the populations of which are characterized by weak adaptive properties, high sensitivity to any change in the habitat or the surrounding areas, slow regeneration after stress and implementation of life strategy in a stable environment (14).

There is a general trend in many parts of Ukraine towards a decrease in the humidity of ecosystems due to prolonged drought, draining of marshes and other wetlands, amelioration of wetlands and their subsequent use for agricultural purposes. As a result, we observe irreversible damage to the ecological balance of ecosystems and depletion in the populations of many medicinal ferns, such as *Adiantum capillus-veneris*, *Marsilea quadrifolia*, *Pilularia globulifera*, *Thelypteris palustris* and others

(Table 1). Further dehumidification of ecosystems threatens their survival. The majority of fern habitats are a fragmented and isolated as a consequence of human activity such as livestock grazing and pastures burning, recreation and tourism because plants cannot cope with the trampling, grazing and waste from stock.

Cutting of forest and unfavorable forest management are further dangers for most of the ferns growing under the canopy of trees, such as *Asplenium adiantum-nigrum*, *A. scolopendrium* L., *Blechnum spicant* (L.) Roth, *Gymnocarpium dryopteris* (L.) Newman, *Onoclea struthiopteris*, *Oreopteris limbosperma* (Bellardi & All.) Holub, *Ophioglossum vulgatum* L. The restoration of populations of these species may last a very long time. There is no information available on current population trends for these species, but the populations are likely to be declining and fragmented following the general trend of destruction and degradation of shallow wetland and forest habitats. In addition to the loss of habitats, overcollection of plants is a significant threat to some rare wild ferns. Many ferns of Ukraine are non-harvest plants, but some of them are used for landscape gardening that may be indicated as a potential risk of extinction.

The state of the majority of the studied medicinal ferns populations is threatened, as they have limited distribution and populations often are represented by isolated individuals and the resources are meager (see Table 1). Thus, harvesting of raw material from the natural environment poses a real threat of extinction.

Analysis of the data on the medicinal effect of biologically active compounds identified in the analyzed ferns showed a wide range of their properties. More than 80% of the ferns have anti-inflammatory, laxative, lactogenic, anthelmintic, diuretic, expectorant, anesthetic, antioxidant and hepatoprotective properties (Table 1). Most of the analyzed ferns (64.1%) have been studied and found to be useful with their anti-inflammatory effect on the stomach and intestines, and their anthelmintic, astringent or light laxative effects. The most famous among them are *Onoclea struthiopteris*, *Polypodium vulgare*, *Pteridium aquilinum* and some others. Concerning diseases of the respiratory system (in-

cluding using them as an expectorant and antipyretic agent), it is known that nearly 45% of the analyzed fern species are used, indicating a vastly researched knowledge about medicinal ferns, which are used to treat respiratory problems. The information about the pharmacological properties or medicinal use of six species (*Asplenium fontanum* (L.) Bernh., *Azolla mexicana* C. Presl, *Cheilanthes acrostica* (Balb.) Tod., *Dryopteris caucasica* (A. Braun) Fraser-Jenk. & Corley, *Dryopteris villarii* (Bellardi) Woyne. ex Schinz & Thell., *Oreopteris limbosperma* (Bellardi & All.) Holun and *Pilularia globulifera* L.) is insignificant to indeterminate, however the main biologically active ingredients were investigated.

The results of the present study demonstrate that these ferns are not as important medicinally as vascular plants in Ukraine, but have considerable importance as obligate components in most of forest systems. Many of them need protection due to irreversible changes in their habitat. The featured table shows the common characteristics of wild medicinal ferns of Ukraine.

## CONCLUSIONS

The use of ferns in Ukraine for their medicinal and other economic value, including ornamental use, is limited, but at the same time, the population densities of many species have decreased primarily due to deforestation and disturbance of habitats. Further research is needed to gather information about the population size, threats and trend of these species throughout their range. Therefore, a significant effort should be taken for the conservation of the species, which are highly sensitive to any change in the habitat. If the protection measures are followed strictly, further studies on ferns as perspective potential for pharmaceuticals could bring many more medicinal benefits.

## REFERENCES

1. Lee CH, Shin SL. Functional Activities of Ferns for Human Health. In: Fernández H, Kumar A, Revilla A, editors. Working with Ferns: Issues and Applications. New York: Springer; 2010. p. 347–359.
2. Manickam VS, Benniamin A. Medicinal Pteridophytes of the Western Ghats, India. Indian Journal of Traditional Knowledge. 2007; 6(4): 611–618.

3. Sarker SK, Hossain ABME. Pteridophytes of greater Mymensingh district of Bangladesh used as vegetables and medicines. *Bangladesh J Plant Taxon.* 2009; 16(1): 47–56.
4. Sharma MP, Mir SA, Mishra AK, Reshi ZA. Preliminary phytochemical screening of some Pteridophyte from district shopian (J & K). *International Journal of Pharmacy and Pharmaceutical Sciences.* 2013; 5(4): 632–637.
5. Valizadeh H, Ali S, Kordi FM, Dehghan H Bahadori MB. Cytotoxicity, Antioxidant Activity and Phenolic Content of Eight Fern Species from North of Iran. *Pharmaceutical Sciences,* 2015; 21: 18–24.
6. Vasudeva SM. Economic importance of Pteridophytes. *SO Indian Fern Journal.* 1999; 16(1–2): 130–152.
7. Plant resources of Russia and neighboring countries: Part I. Families Lycopodiaceae – Ephedraceae; Part II. Additions to the 1-7th volumes. Budantsev AL editor. SPb: Peace and Family-95; 1996 (In Russian).
8. Minarchenko VM. Medical vascular plants of Ukraine (medical and resource value). Kyiv: Fitosociocentre; 2005 (In Ukrainian).
9. Han X, Li Z, Li CY, Jiaa WN, Wang HT. Phytochemical Constituents and Biological Activities of Plants from the Genus *Dryopteris*. *Chemistry & Biodiversity.* 2015; 12: 1131–1162.
10. De Medeiros JM, Macedo M, Contancia JP, Nguyen C, Cunningham G, Miles DH. Antithrombin activity of medicinal plants of the Azores. *J Ethnopharmac.* 2000; 72 (1–2): 157–165.
11. Vasheka OV, Bezsmertna OO. Ferns atlas of Ukrainian flora: monograf. Kyiv: Palyvoda A.V.; 2012 (In Ukrainian).
12. Didukh Ya.P. (ed). *Red Book of Ukraine.* Kyiv: Globalkonsaltyng, 2009.
13. The IUCN Red List of Threatened Species [Internet]. Version 2016-3. [Downloaded on 2017 January 24; cited 2017 March 02]. Inc. Available from: <http://www.iucnredlist.org>
14. Minarchenko V. Medicinal plants of Ukraine: diversity, resources, legislation. *Medicinal Plant Conservation Newsletter.* 2011; 14: 7-13. [http://cmsdata.iucn.org/downloads/mpc\\_14\\_1.pdf](http://cmsdata.iucn.org/downloads/mpc_14_1.pdf)
15. Al-Snafi AE. The chemical constituents and pharmacological effects of *Adiantum capillus-veneris* – a review. *Asian Journal of Pharmaceutical Science & Technology.* 2015; 5(2): 106–111.
16. Singha M, Singha N, Khareb PB, Rawata AKS. Antimicrobial activity of some important *Adiantum* species used traditionally in indigenous systems of medicine. *J Ethnopharmac.* 2008; 115(2): 327–329.
17. Ishaq MS, Hussain MM, Siddique M, Ali G, Khatkhat M, Ahmad S, et al. In Vitro Phytochemical, Antibacterial, and Antifungal Activities of Leaf, Stem, and Root Extracts of *Adiantum capillus veneris*. *The Scientific World Journal.* 2014. doi. [org/10.1155/2014/269793](http://dx.doi.org/10.1155/2014/269793).
18. Sen A, Gosh PD. A note on the ethnobotanical studies of some pteridophytes in Assam. *Indian J Tradit Knowl.* 2011; 10: 292–295.
19. Zedan Z I, Ahmed A S, Gouda Y G. Phytochemical and biological studies of *Adiantum capillus-veneris* L. *Saudi Pharmaceutical Journal.* 2011; 19(2): 65–74.
20. Mannar Mannan M, Maridass M, Victor B. A Review on the Potential Uses of Ferns. *Ethnobotanical Leaflets.* 2008; 12: 281–285.
21. Dar A, Faizi S, Naqvi S, Roome T, Zikr-ur-Rehman S, Ali M, et al. Analgesic and antioxidant activity of mangiferin and its derivatives: the structure activity relationship. *Biol Pharm Bull.* 2005; 28(4): 596–600.
22. Dall'Acqua S, Tomè F, Vitalini S, Agradi E, Innocenti G. In vitro estrogenic activity of *Asplenium trichomanes* L. extracts and isolated compounds. *J Ethnopharmacol.* 2009; 122(3): 424–429.
23. Imperato F. A new sulfated flavonol glycoside in the fern *Asplenium fontanum*. *Bernh Chem Ind.* 1980; 540–541.
24. Imperato F. Xanthone 2,4-di-C-glycosides from *Asplenium adiantum-nigrum*. *Phytochemistry.* 1991; 30: 3839–3840.
25. Iwashina T, López-Sáez JA, Herrero A, Kitajima J, Matsumoto S. Flavone glycosides from *Asplenium foreziense* and its five related taxa and *A. incisum*. *Biochemical Systematics and Ecology.* 2000; 28: 665–671.
26. Fan P, Zhao L, Hostettmann K, Lou H. Chemical constituents of *Asplenium ruta-muraria* L. *Natural Product Research.* 2012; 26(15): 1413–1418.
27. Iwashina T, Matsumoto S. Flavonoid Properties of six *Asplenium* species in Vanuatu and New Caledonia, and Distribution of Flavonoid and Related

- Compounds in *Asplenium*. Bull Natl Mus Nat Sci. 2011; Ser. B, 37(3): 133–145.
28. Ashalatha K. Isolation & characterization of biologically active compounds from the medicinal plants *Salvinia natans* and *Decalepis hamiltonii* and total synthesis of cleistenolide and yashabushidiol derivatives. 2011 [updated 2013 Apr. 22; cited 2017 Jan 20]. Available from: <http://hdl.handle.net/10603/8226>
29. Mizuno M, Kyotani Y, Iinuma M, Tanaka T, Kojima H, Iwatsuki K. Kaempferol glycosides in *Asplenium scolopendrium* Newm. Zeitschrift für Naturforschung C. 1990; 45: 143–146.
30. You-Min S, Young-Won C, Min-Hye Y, Jin-Woong K. Terpenoid constituents from the aerial parts of *Asplenium scolopendrium*. Natural Product Sciences. 2008; 14(4): 265–268.
31. Aibuldinov Ye, Ozek G, Rakhmadiyeva S, Bashier KHC. Gas-chromatographic and mass spectrometric evaluation of essential oils of four ferns from Kazakhstan. Newsletter of L.N.Gumilyov Eurasian National University. 2012; 6: 265–270.
32. Imperato F. Flavonol glycosides from ferns of the genera *Asplenium* and *Cheilanthes*. Biochemical Systematics and Ecology. 1989; 17: 161–166.
33. Froissard D, Fons F, Bessière JM, Buatois B, Rapior S. Volatiles of French ferns and «fougère» scent in perfumery. Nat Prod Commun. 2011; 6(11): 1723–1726.
34. Singh S, Singh R. Ethnomedicinal use of Pteridophytes in reproductive health of tribal women of Pachmarhi biosphere reserve, Madhya Pradesh, India. International Journal of Pharmaceutical Sciences and Research. 2012; 3(12): 4780–4790.
35. Iwashina T, Matsumoto S. Flavonoid variation and evolution in *Asplenium normale* and related species (*Aspleniaceae*). Journal of Plant Research. 1994; 107(3): 275–282.
36. Akmuradov A, Rakhmanov O. Medicinal resources of pteridoflora of Turkmenistan, applied to folk medicine. Healthcare of Turkmenistan. 2011; 1: 29–34.
37. Akmuradov A, Shaiymov BK. Medical plants of the Koitendag flora used in folk medicine. Siberian Medical Journal. 2015; 4: 86–89.
38. Bahadori MB, Kordi FM, Ahmadi AA., Bahadori Sh, Valizadeh H. Antibacterial evaluation and preliminary phytochemical screening of selected ferns from Iran. Research Journal of Pharmacognosy (RJP). 2015; 2(2): 53–59.
39. Kumar A, Kumari J, Kumar H, Nath A., Singh JK., Ali M, et al. Hepatoprotective and antioxidant effect of *Azolla filiculoides* on profenofos induced hepatotoxicity in swiss albino mice. Carib J Sci Tech. 2014; 2: 372–377.
40. Pieterse AH, de Lange L, Van Vliet JP. A comparative study of *Azolla* in the Netherlands. Acta Botanica Neerlandica. 1977; 26(6): 433–449.
41. Pereira AL, Carrapiço F. Histochemistry of simple hairs from the foliar cavities of *Azolla filiculoides*. Plant Biosystems. 2007; 141(3): 323–328.
42. Masood A, Zeeshan M, Abraham G. Response of growth and antioxidant enzymes in *Azolla* plants (*Azolla pinnata* and *Azolla filiculoides*) exposed to UV-B. Acta Biol Hung. 2008; 59 (2):247-258.
43. Holst RW. Anthocyanins of *Azolla*. American Fern Journal. 1977; 67: 99–100.
44. Plants for a future [Internet]. [updated 2014 Oct 27; cited 2017 May 02]. Available from: <http://www.pfaf.org/user/Default.aspx>
45. Srivastava K. Ethnobotanical Studies of Some Important Ferns. Ethnobotanical Leaflets. 2007; 11: 164–172.
46. Đurdevih L, Mitrović M, Pavlović P, Bojović S, Jarić S, Oberan L, et al. Total phenolics and phenolic acids content in leaves, rhizomes and rhizosphere soil under *Ceterach officinarum* D.C., *Asplenium trichomanes* L. and *Adiantum nigrum* L. in the gorge of Sićevo (Serbia). Ekológia (Bratislava). 2007; 26(2): 164–173.
47. Guarino C, De Simone L, Santoro S. Ethnobotanical Study of the Sannio Area, Campania, Southern Italy. Ethnobotany Research & Applications. 2008; 6: 255–317.
48. Ullah F, Ullah A, Sohail A, Khan R. Diversity, distribution and ecological importance of weeds in the maize crop at maidan valley, dir (I), khyber pakh-tunkhwa, Pakistan. Pak J Weed Sci Res. 2015; 21(4): 543–553.
49. Vogler G, Donath O, Saukel J, Rauch AW, Kählig H, Krenn L. Polar phenolic compounds in *Dryopteris filix-mas* and *Dryopteris dilatata*. Verh Zool-Bot Ges Österreich. 2012; 148/149: 279–289.
50. Imperato F. Kaempferol 3-O-(acetylrutinoside), a new flavonoid and two new fern constituents, quercetin 3-O-(acetylglucoside) and 3-O-

- (acetylrutinoside) from *Dryopteris villarii*. American Fern Journal. 2006; 96(3): 93–96.
51. Kovganko NV, Kashkan ZhN, Krivenok SN, Potapovich MV, Eremin AN, Metelitsa DI. Bioactive compounds in the flora of Belarus. 2. Astragalins, an effective protector of catalase from ultrasonic inactivation in aqueous solutions. Chemistry of Natural Compounds. 2004; 40(1): 71–74.
  52. Uma R, Pravin B. In vitro Cytotoxic Activity of *Marsilea quadrifolia* Linn of MCF-7 Cells of Human Breast Cancer. Int Res J Medical Sci I. 2013; 1(1): 10–13.
  53. Ripa FA, Nahar L, Haque M, Islam MM. Antibacterial, Cytotoxic and Antioxidant Activity of Crude Extract of *Marsilea quadrifolia*. European Journal of Scientific Research. 2009; 33(1): 123–129.
  54. Li S, Zhang D, Yang L, Li Y, Zhu X, Kmonílková E, et al. HPLC Quantitative Analysis of Main Stilbenes and Flavones in Different Parts of *Matteuccia struthiopteris*. Journal of Chemistry. 2013; 2013. doi.org/10.1155/2013/452610
  55. Li B, Ni Y, Zhu LJ, Wu FB, Yan F, Zhang X, et al. Flavonoids from *Matteuccia struthiopteris* and Their Anti-influenza virus (H1N1) activity. J Nat Prod. 2015; 78(5): 987–95.
  56. Zhang D, Yang L, Fu MH, Tu YY. Studies on chemical constituents of rhizome of *Matteuccia struthiopteris* III. China Journal of Chinese Materia Medica. 2008; 33(14): 1703–1705.
  57. Yang L, Wang MY, Zhao YY, Tu YY. Chemical constituents of the rhizome of *Matteuccia struthiopteris*. Acta Pharmaceutica Sinica. 2005; 40(3): 252–254.
  58. Yang L, Wang MY, Zhao YY, Tu YY. Studies on chemical constituents in rhizome of *Matteuccia struthiopteris*. China Journal of Chinese Materia Medica. 2004; 29(7): 648–649.
  59. Yang L, Zhao YY, Tu YY. Chemical constituents of the rhizome of *Matteuccia struthiopteris*. China Journal of Chinese Materia Medica. 2003; 28: 278–279.
  60. Boroznets AS, Andrianova GS. The search for carriers of vitamins P and C among wild flora of the Far East. Proceedings of the Khabarovsk Medical Institute. 1962; 23(1): 147–150 (In Russian).
  61. Shreter AI. Medicinal flora of the Soviet Far East. Moscow: Medicine; 1975 (In Russian).
  62. Clericuzio M, Burlando B, Gandini G, Tinello S, Ranzato E, Martinotti S, et al. Keratinocyte wound healing activity of galactoglycerolipids from the fern *Ophioglossum vulgatum* L. J Nat Med. 2014; 68(1): 31–37.
  63. Clericuzio M, Tinello S, Burlando B, Ranzato E, Martinotti S, Cornara L, et al. Flavonoid oligoglycosides from *Ophioglossum vulgatum* L. having wound healing properties. Planta Med. 2012; 78(15): 1639–1644.
  64. Fons F, Froissard D, Bessière JM, Buatois B, Rapior S. Biodiversity of Volatile Organic Compounds from Five French Ferns. Nat Prod Commun. 2010; 5(10): 1655–1658.
  65. Imperato F. A new flavonoid, quercetin 3-O-(X''-acetyl-X''-cinnamoyl-glucoside) and a new fern constituent, quercetin 3-O-(glucosylrhamnoside) from *Dryopteris villarii*. American Fern Journal. 2007; 97: 233–236.
  66. Assessment report on *Polypodium vulgare* L., rhizoma. European Medicines Agency Evaluation of Medicines for Human Use. Doc. Ref.: EMEA/HMPC/600669/2007 [Internet]. 2008. [updated 2008 Nov 6; cited 2017 May 25]. Available from: [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Herbal\\_-HMPC\\_assessment\\_report/2009/12/WC500018581.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Herbal_-HMPC_assessment_report/2009/12/WC500018581.pdf)
  67. Soare LC, Ferdeş M, Deliu I, Gibea A. Studies regarding the antibacterial activity of some extracts of native pteridophytes. UPB Sci Bull. 2012; Series B, 74(1): 21–26.
  68. Shishmarev VM. Medical ferns of Burytia. Buryat State University Bulletin. 2013; 12: 69–71 (In Russian).
  69. Chen N. Study on the extraction of flavonoid compound in *Pteridium aquilinum* (L.) and its antioxidant property. Food and Ferment Indust. 2003; 29(11): 63–66.
  70. Xu W, Zhang F, Luo Y, Ma L, Kou X, Huang K. Antioxidant activity of a water-soluble polysaccharide purified from *Pteridium aquilinum*. Carbohydr Res. 2009; 344(2): 217–222.
  71. Srilaxmi P, Sareddy GR, Kavikishor PB, Setty OH, Babu PP. Protective efficacy of natansnin, a dibenzoyl glycoside from *Salvinia natans* against CCl<sub>4</sub> induced oxidative stress and cellular degeneration in rat liver. BMC Pharmacology. 2010; 10:13. DOI: 10.1186/1471-2210-10-13

72. Yang X et al. Encyclopedic reference of traditional Chinese medicine. Berlin Heidelberg: Springer-Verlag; 2003.
73. Murakami T, Wada H, Tanaka N, Chen CM. Chemical and Chemotaxonomical Studies of Filices. LXV.: A Few New Flavonoid Glycosides. (2) (Organic). Journal of the Pharmaceutical Society of Japan. 1986; 106(11): 982–988.